

# Kinetics Analysis of Synthesis Reaction of Struvite With AirFlow Continous Vertical Reactors

*by* Luluk Edahwati

---

**Submission date:** 26-Nov-2020 09:18AM (UTC+0700)

**Submission ID:** 1457308361

**File name:** 7.3.Kinetics\_Analysis\_of\_Synthesis\_Reaction.pdf (695.77K)

**Word count:** 2720

**Character count:** 14254

PAPER • OPEN ACCESS


2

## Kinetics Analysis of Synthesis Reaction of Struvite With Air-Flow Continous Vertical Reactors

1

To cite this article: L Edahwati *et al* 2018 *J. Phys.: Conf. Ser.* **953** 012245

View the [article online](#) for updates and enhancements.



**IOP ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

## <sup>2</sup> Kinetics Analysis of Synthesis Reaction of Struvite With Air-Flow Continuous Vertical Reactors

L Edahwati, S Sutiyono<sup>1</sup>, S Muryanto<sup>2</sup>, J Jamari<sup>3</sup> dan A P Bayuseno<sup>3</sup>

<sup>1</sup>Universitas Pembangunan Nasional "Veteran" Jawa Timur, Surabaya

<sup>2</sup>Department of Chemical Engineering, UNTAG University in Semarang.

<sup>3</sup>Department of Mechanical Engineering, Diponegoro University, Semarang

<sup>1</sup>edahwatiluluk@yahoo.co.id

\*apbayuseno@gmail.com

**Abstract.** Kinetics reaction is a knowledge about a rate of chemical reaction. The differential of the reaction rate can be determined from the reactant material or the formed material. The reaction mechanism of a reactor may include a stage of reaction occurring sequentially during the process of converting the reactants into products. In the determination of reaction kinetics, the order of reaction and the rate constant reaction must be recognized. This study was carried out using air as a stirrer as a medium in the vertical reactor for crystallization of struvite. Stirring is one of the important aspects in struvite crystallization process. Struvite crystals or magnesium ammonium phosphate hexahydrates ( $MgNH_4PO_4 \cdot 6H_2O$ ) is commonly formed in reversible reactions and can be generated as an orthorhombic crystal. Air is selected as a stirrer on the existing flow pattern in the reactor determining the reaction kinetics of the crystal from the solution. The experimental study was conducted by mixing an equimolar solution of 0.03 M  $NH_4OH$ ,  $MgCl_2$  and  $H_3PO_4$  with a ratio of 1: 1: 1. The crystallization process of the mixed solution was observed in an inside reactor at the flow rate ranges of 16-38 ml/min and the temperature of 30°C was selected in the study. The air inlet rate was kept constant at 0.25 liters/min. The pH solution was adjusted to be 8, 9 and 10 by dropping wisely of 1 N KOH solution. The crystallization kinetics was examined until the steady state of the reaction was reached. The precipitates were filtered and dried at a temperature for subsequent material characterization, including Scanning Electron Microscope (SEM) and XRD (X-Ray diffraction) method. The results show that higher flow rate leads to less mass of struvite.

### 1. Introduction

<sup>2</sup> Kinetics reaction is the branch of chemistry that studies about chemical reaction. The differential of the reaction rate can be determined from the reactant material or the formed material. The reaction mechanism of a reactor may include a stage of reaction occurring sequentially during the process of converting the reactants into products. Meanwhile, the rate of chemical reaction is the number of moles of reactants per unit volume in units of time [15]. The kinetic equation of the reaction is divided into two equations ie the kinetic velocity of the homogeneous reaction and the heterogeneous. In this study, the researchers determined the kinetic's data using struvite solution. In our body can also form struvite that can harm us. In the body, struvite can be formed naturally in the process of formation of urine. Even struvite that settles in our bodies can lead to a disease that kidney stone disease [5].

According to [112], the experimental observations for struvite precipitation kinetics with different stirrer speeds clearly indicate that the mixing intensity used in this study has an influence on the reaction kinetics constants. This study using the vertical reactor as a crystalline struvite crystal formation's medium. The crystalline struvite in the present study was formed from synthetic waste comprising  $MgCl_2$ ,  $NH_4OH$ ,  $H_3PO_4$  and using KOH as a pH. Researchers tried to find the kinetic of reaction struvite crystals and calculate the reaction constant in the continuous process of crystallization struvite by using the vertical reactor. The precipitates were filtered and dried at a temperature for subsequent material characterization, including Scanning Electron Microscope (SEM) and XRD (X-Ray diffraction) method. The results show that higher flow rate leads to less mass of struvite.

The process of struvite crystals formation by using the crystallization process. Crystallization is the process by which solid particles are formed from the homogeneous phase [4]. The formation of these struvite crystals uses a continuous process. In the previous research [14], struvite forming process has been carried out using continuous flow stirred tank reactor and after Batch Process is done, the result is better than batch process. Then According to previous researchers [7],  $Mg^{2+}$  + molar ratio factor:  $NH_4^+$ :  $PO_4^{3-}$  influenced in the experiment and speed gradient. According to [12] with different Mg: P ratios indicating that the higher the ratio (in the range 1.0-1.6), the better the struvite formation efficiency. So the kinetics precipitation also influential. Then according to [13], the reaction rate decreases with increasing precipitation time with minimum phosphorus and magnesium concentrations. Kinetics of the struvite crystallization process are influenced by temperature. The struvite solubility product increases as the temperature increases. Moreover, an increase in the reaction temperature causes an increase in the reaction rate parameter. Besides, the kinetics of the struvite reaction are influenced by the pH of the aqueous solution because the availability of struvite ions is affected by pH. [2].

### 1.1 Type Of Kinetics Reaction

#### 1. Homogeneous Reaction

A homogeneous reaction is a reaction involving a phase reaction mixture. The reaction velocity can be expressed as the volume of the reaction mixture, so mathematically the reaction rate can be written as follows :

$$-r_A = 1/V = dNA/dt \quad (1)$$

In addition to the speed of the reaction the whole material becomes :

$$\frac{-r_A}{a} = \frac{-r_B}{b} = \frac{-r_R}{r} = \frac{-r_S}{s} \quad (2)$$

#### 2. Heterogeneous Reactions

Heterogeneous reactions are reactions that involve multiple phases in the reaction. [18].

### 1.2 Residence Time Of Reactor With Continuous Process

In a continuous process, known as residence time. According to [1]. The theory of residence time relates to how the incoming particle flows through it and leaves a system. It intuitively looks reasonable to expect that not all particles have the same residence time. For constant density, levenspiel shows that:

$$\tau = Vol/Q \quad (3)$$

where :


$\tau$  = Residence Time (minute)

V = Volume (cc)

Q = Debit (cc/minute)

### 1.3 Crystallisation of Struvite

Struvite is magnesium Ammonium phosphate ( $MgNH_4PO_4 \cdot 6H_2O$ ) which usually crystallizes in reversible reactions and produces orthorhombic crystals with  $Mg^{2+}$  :  $NH_4^+$  :  $PO_4^{3-}$ . [2]. The reaction of struvite formation may be expressed as follows :



Struvite occurs spontaneously in solution when the content in  $Mg^{2+}$ ,  $NH_4^+$  and  $PO_4^{3-}$  reaches a 1:1:1 molar ratio under specific conditions of pH, temperature, mixing energy and presence of foreign

ions [3]. Although conditions affecting success of P removal and recovery as struvite and their consequences on thermodynamics of precipitation have been widely investigated [11][9], knowledge on kinetics of struvite precipitation, that is to say on rates at which struvite forms, is still limited [10]. Struvite crystallisation can be divided in two major steps: nucleation and crystal growth [6].

## 2. Method

In this study, researchers use a vertical reactor as a medium for the formation of struvite crystals. Struvite crystals or magnesium ammonium phosphate hexahydrates ( $MgNH_4PO_4 \cdot 6H_2O$ ) is commonly formed in reversible reactions and can be generated as an orthorhombic crystal.




Figure.1. Vertical Reactor

Air is selected as a stirrer on the existing flow pattern in the reactor determining the reaction kinetics of the crystal from the solution. The first thing to do is make the Magnesium Ammonium Phosphat (MAP) with the concentration ratio  $Mg: NH_4: PO_4 = 1: 1: 1$  and KOH 1N solution. Then compile set of tools as attached in Fig. 1. Then fill the reactor vertical with MAP reached the  $\frac{3}{4}$  height of the reactor and KOH to control the pH of the solution. Turn on the stove with temperature setting  $30^\circ C$ . Set the incoming MAP rate of 15,20,25,30,40 ml / min and KOH rate. Set the MAP and KOH rates in 1: 2 ratio. Turn on the compressor to drain the air into the reactor, set the air rate to 0.25 ml / min. Collect the mixture of MAP and KOH and measure the pH of mixed MAP and KOH solution to reach pH 8,9,10 and wait until 5 minutes. Then collect the result and filter using filter paper. Dry the struvite's crystal and record the weight of the precipitate. Analyze the sediment.

## 3. Result and Discussion


### 3.1 Determine the Order Reaction

The observed variation pH for crystallization struvite during the study was given for pH 8, 9, and 10. The study was carried out at a temperature of  $30^\circ C$ . After analyzing the struvite crystals, the final concentration of struvite (Ca) is obtained as shown in Table 1. With the variation of pH, the conversion price (Xa) increases with the increase of pH. Then, with the data so that it can be determined the determination of the reactions. it can be seen that the determination of reaction kinetics in this struvite crystal leads to first order.


**Table 1.** Calculation Order Reaction

pH	$\tau$ Residence Time (minute)	Temperature (T) 30 <sup>o</sup> C			
		C <sub>A0</sub> (%)	C <sub>A</sub> (%)	X <sub>a</sub>	-ln (1-X <sub>a</sub> )
8	33.25	0.43609	0.41928	0.007667	0.00769615
	24.3975	0.43609	0.42271	0.01591	0.01603794
	19.95	0.43609	0.42291	0.016391	0.0165265
	16.625	0.43609	0.423	0.016607	0.01674643
	12.46875	0.43609	0.42395	0.01889	0.01907084
9	33.25	0.43609	0.4029	0.0317	0.03221319
	24.3975	0.43609	0.403	0.03146	0.03196502
	19.95	0.43609	0.4031	0.031219	0.03171691
	16.625	0.43609	0.4032	0.030979	0.03146887
	12.46875	0.43609	0.40325	0.030859	0.03134487
10	33.25	0.43609	0.37912	0.088851	0.09304881
	24.3975	0.43609	0.381	0.084333	0.08810221
	19.95	0.43609	0.38275	0.080127	0.08351955
	16.625	0.43609	0.385	0.074719	0.07765825
	12.46875	0.43609	0.38711	0.069648	0.07219269


3.2 Calculated of Order Reaction and Result Scanning electron micrographs (SEM) of struvite



**Figure 2.** Graphic  $\tau$  (Residence Time) vs  $-\ln(1-X_a)$  and Result Scanning electron micrographs (SEM) of struvite at the pH of 8



**Figure 3.** Graphic  $\tau$  (Residence Time) vs  $-\ln(1-X_a)$  and Result Scanning electron micrographs (SEM) of struvite at the pH of 9



**Figure 4.** Graphic  $\tau$  (Residence Time) vs  $-\ln(1-X_a)$  and Result Scanning electron micrographs (SEM) of struvite at the pH of 10

The graphic between  $\tau$  (residence time) and  $-\ln(1-X_a)$  shows a straight graph containing  $R^2$  which the value close to 1 as shown in fig.2, fig.3, and fig.4. This proves that the kinetics of the struvite synthesis reaction follow the order first and with the variation of pH (8,9,10).

**Table 2.** Calculation % Error

pH	R <sup>2</sup>	R	Y = k	t	+	1/C <sub>A0</sub>	% Kesalahan R
8	0.87625	0.93608	y = 0.0005	x	+	0.02131	6.39%
9	0.96775	0.98374	y = 0.00004	x	+	0.07775	1.62%
10	0.94358	0.97138	y = 0.00101	x	+	0.10838	2.86%

While in table 2 we can be obtained data rate reaction constant at each pH is for pH 8 = 0.0005; pH 9 = 0,00004; and pH 10 = 0,00101 and for pH 8, pH 9, pH 10 got percentage of error respectively that is 6,39%; 1.62%; 2.86%. So that has the smallest percent error is at pH 9, where pH 9 is the best conditions for struvite formation.

#### 4. Conclusion

After plotting between  $\tau$  (residence time) and  $-\ln(1-X_a)$  at the 1st reaction order, a straight graph containing R<sup>2</sup> values approaches 1, this proves that the kinetics of the struvite synthesis reaction follow the first order and with the pH variation (8,9,10) obtained the price of% mistake consecutively is 6.39%; 1.62%; 2.86%. it can be concluded that at pH 9% obtained the smallest error that is equal to 1.62%, so this condition is the best condition in the process of crystalline struvite formation. While the reaction rate constant obtained at the first order of reaction at pH variation (8,9,10) was 0.5.10<sup>-3</sup>, respectively; 0.04.10<sup>-3</sup>; 1.01.10<sup>-3</sup> min<sup>-1</sup>. The best results of struvite crystals formed at pH 9. Based on the results of scanning electron micrographs (SEM) analysis at pH 9 obtained a struvite content of 27.85%

#### 5. Acknowledgment

The author wish to thank to University of Pembangunan Nasional "Veteran" Surabaya for supporting this research programs.

#### 6. References

- [1] Ahmmad Shukrie Bin Md Yudin.2012.*Studies On The Residence Time Distribution Of Solids In A Swirling Fluidized Bed*, Perak: Thesis of Mechanical Engineering,p.20-21.
- [2] Crutchik, D. 2016. *Kinetics of the reversible reaction of struvite crystallization*. Department of Chemical Engineering, Institute of Technology, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain
- [3] Doyle J.D. and Parsons S.A., Struvite formation, control and recovery. *Water Res.*, 36, 3925-3940, (2002).
- [4] Geankoplis, Christie J. 1983. *Transport Processes and Unit Operation*. America: Prentice Hall International.inc.
- [5] Hao, X,-D. 2008. *Struvite formation, analytical methods and effects of pH and Ca<sup>2+</sup>*. *Journal of Civil Engineering*
- [6] Jones A.G., Crystallization process system. Butterworth/ Heinemann, UK, (2002).
- [7] Kumar, Ramesh, Parimal Pal. 2013. *Turning hazardous waste into value-added products: production and characterization of struvite from ammoniacal waste with new approaches*. Environment and Membrane Technology Laboratory, Department of Chemical Engineering, National Institute of Technology, Durgapur 713209, India
- [8] Levenspiel.1999.*Chemical Reaction Engineering*.New York:John Wiley & Sons
- [9] Michalowski T. and Pietrzyk A., A thermodynamic study of struvite + water system. *Talanta*,68, 594-601, (2006).

- [10] Nelson N.O., Mikkelsen R.L. and Hesterberg D.L., Struvite precipitation in anaerobic swine lagoon liquid: effect of pH and Mg:P ratio and determination of rate constant. *Bioresource Technol.*, 89, 229-236, (2003).
- [11] Ohlinger K.N., Young T.M. and Schroeder E.D., Predicting struvite formation in digestion. *Water Res.*, 32, 3607-3614, (1998).
- [12] Rahaman, M.S, 2003. *Effects of Various Process Parameters on Struvite Precipitation Kinetics and Subsequent Determination of Rate Constants*. Canada: Journal of Civil Engineering Vol. 3, p.2
- [13] Shalaby, Marwa Saied and Shadia El-Rafie. 2015. *Struvite Precipitation and Phosphorous Removal from Urine Synthetic Solution: Reaction Kinetic Study*. Chemical Engineering and Pilot Plant Department, National Research Center, El buhouth St., Dokki, Giza 12311, Egypt
- [14] Warmadewanthi, J.C. Liu. 2009. *Recovery of phosphate and ammonium as struvite from semiconductor wastewater*. Department of Chemical Engineering, National Taiwan University of Science and Technology, 43 Keelung Road, Section 4, Taipei 106, Taiwan
- [15] Widodo and Maesroh:(2016). *Kinetics Study of Methylacetate's Reaction From Acetic Acid and Methanol with Variable Time, Catalyst Concentration and Comparison of Reactants*. Journal of Chemical Engineering Vol. 3,p.2



# Kinetics Analysis of Synthesis Reaction of Struvite With AirFlow Continous Vertical Reactors

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

4%

PUBLICATIONS

12%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Politeknik Negeri Bandung Student Paper	6%
2	worldwidescience.org Internet Source	4%
3	dspace.lib.cranfield.ac.uk Internet Source	4%
4	D. Crutchik, J.M. Garrido. "Kinetics of the reversible reaction of struvite crystallisation", Chemosphere, 2016 Publication	2%
5	re.public.polimi.it Internet Source	2%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Kinetics Analysis of Synthesis Reaction of Struvite With AirFlow Continous Vertical Reactors

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---