



Manuscript Information**Authors****Files**
 **Download PDF Version of Manuscript**

#	File Type	File Name	Size	File Description	Upload Date
 Files Sent by Authors					
1	Title Page	Title Page.docx	12.9 KB		2019-09-30
2	Research Highlights	HIGHLIGHTS.docx	13.75 KB		2019-09-30
3	Graphical Abstract	GRAPHICAL ABSTRACT.docx	239.17 KB		2019-11-29
4	Manuscript Main File	Revised_2352 PR3.doc	1.86 MB		2019-11-30
5	Manuscript Main File (Changes Highlighted)	Revised_2352 PR3.doc	1.86 MB		2019-11-30
6	Response to Reviewer	Author Query Form.doc	60.5 KB		2019-11-30
 Files Sent by Editor-in-Chief to Author					
7	Editor-in-Chief File	2352 PR3.doc	1.83 MB		2019-11-29
 Files Sent by Author (Proof)					
8	Copyright Transfer Agreement	Copyright-1.pdf	64.14 KB		2019-11-30
9	Conflict of Interest Disclosure Form	Conflict of Interest.pdf	63.44 KB		2019-11-30
10	Galley Proof	2352 Galley Proof.doc	1.86 MB		2019-11-30



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acknowledgement of Submission (Manuscript #GJESM-1909-2352)

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir>

Tue, Oct 8, 2019 at 1:57 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com, nourijafar@gmail.com

Manuscript ID: GJESM-1909-2352

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

I wish to acknowledge receiving of the above-mentioned manuscript. It should be noted that the manuscript will be reviewed for possible publication in the **Global Journal of Environmental Science and Management (GJESM)**.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Upon submission, the manuscript is immediately screened for similarity through trustworthy software named iThenticate to be assured about its originality.

Our editorial decision will be brought to your attention once the paper has been reviewed due to the referees' consideration. Please also note that manuscript submission, processing, and publication are free of charge in GJESM Journal.

It is necessary for you, as the corresponding author of the submitted manuscript, to declare that you have informed all the co-author(s) of the manuscript of making this submission and you have obtained their permission to proceed with it on their behalf.

Thank you for your interest in contributing to the **Global Journal of Environmental Science and Management** and choosing it as your research hub.

Truly yours,

Professor J. Nouri

Editor in Chief

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Manuscript Needs Resubmission (Manuscript #GJESM-1909-2352)

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Mon, Sep 30, 2019 at 10:26 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com

Dear Author

1- Attached, through the system on your dashboard, you will find your modified manuscript (#2352) according to the GJESM Journal style. Thus, in order you could get more positive feedback of the manuscript peer reviewing, you have to revise your manuscript technically and in English language revising as quickly as possible according to the provided comments which are indicated on the current manuscript text. Thus, do not change the modified text style.

2- Remember correct your further revisions JUST on the prepared current manuscript file which is attached into your portal in the system, not at your previous submitted file. Besides, all of your amendments must be highlighted with RED fonts color to be recognized by Editor. Afterwards, the manuscript revision must be returned back by at most 10 days not more for the further processing to see if you are serious your manuscript to be processed.

3- After the deadline, in case of not receiving your revised manuscript, it will be archived with no further action. Upon completion, the revised manuscript must be re-submitted through the website system on your own portal back where you had already submitted your manuscript not occupying a new ID number.

4- Finally, GJESM Journal as free charge publication is indexed in Web of Science and Scopus, tries to end a submitted manuscript peer reviewing within a month or less for the possible manuscript publication. Thus, you have to revise your manuscript as carefully and quickly as possible according to the provided comments which are outlined on the current manuscript text.

Your delay in resubmitting will lead to archiving the manuscript with no further action.

Editorial Office



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Editorial system registration

1 message

Global Journal of Environmental Science and Management

Mon, Sep 30, 2019 at 6:40

<journal@iranjournals.ir>

PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Dear Euis Nurul Hidayah,

Welcome to **Global Journal of Environmental Science and Management** website.

Thank you for your registration in the Editorial System Online Submission and peer review tracking system.
Below please find your username and confidential password, which you need to access the system at:

<https://www.gjesm.net/>

Your username is: euisnh@gmail.com

Your password is: **1ga7bu9khg**

Please save this information in a safe place.

Once you login , you can change your password on your profile.

Editorial Office

Global Journal of Environmental Science and Management

<https://www.gjesm.net/>



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acknowledgement of Submission (Manuscript #GJESM-1909-2352)

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Mon, Sep 30, 2019 at 7:22 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com, nourijafar@gmail.com

Manuscript ID: GJESM-1909-2352

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

I wish to acknowledge receiving of the above-mentioned manuscript. It should be noted that the manuscript will be reviewed for possible publication in the **Global Journal of Environmental Science and Management (GJESM)**.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Upon submission, the manuscript is immediately screened for similarity through trustworthy software named iThenticate to be assured about its originality.

Our editorial decision will be brought to your attention once the paper has been reviewed due to the referees' consideration. Please also note that manuscript submission, processing, and publication are free of charge in GJESM Journal.

It is necessary for you, as the corresponding author of the submitted manuscript, to declare that you have informed all the co-author(s) of the manuscript of making this submission and you have obtained their permission to proceed with it on their behalf.

Thank you for your interest in contributing to the **Global Journal of Environmental Science and Management** and choosing it as your research hub.

Truly yours,

Professor J. Nouri

Editor in Chief

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acknowledgement of Submission (Manuscript #GJESM-1909-2352)

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Wed, Oct 9, 2019 at 8:16 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com, nourijafar@gmail.com

Manuscript ID: GJESM-1909-2352

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

I wish to acknowledge receiving of the above-mentioned manuscript. It should be noted that the manuscript will be reviewed for possible publication in the **Global Journal of Environmental Science and Management (GJESM)**.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Upon submission, the manuscript is immediately screened for similarity through trustworthy software named iThenticate to be assured about its originality.

Our editorial decision will be brought to your attention once the paper has been reviewed due to the referees' consideration. Please also note that manuscript submission, processing, and publication are free of charge in GJESM Journal.

It is necessary for you, as the corresponding author of the submitted manuscript, to declare that you have informed all the co-author(s) of the manuscript of making this submission and you have obtained their permission to proceed with it on their behalf.

Thank you for your interest in contributing to the **Global Journal of Environmental Science and Management** and choosing it as your research hub.

Truly yours,

Professor J. Nouri

Editor in Chief

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Manuscript Authors Contributions

2 messages

GJESM Journal <gjesm.publication@gmail.com>

Tue, Dec 31, 2019 at 6:16 PM

To: Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

(URGENT PLEASE)!

Dear Author(s),

In order to finalize your article for the entire publication, the AUTHORS CONTRIBUTIONS item must be included for all accepted articles in the journal. Thus, please write the contribution of each authors regarding your study into a few lines and send us via email immediately to be included in your current article publication.

Your delay may lead to be postponed your entire publication

Truly yours

Editorial Office

Global Journal of Environmental Science and Management (GJESM)

Tel.: +9821- 26105110

Fax: +9821- 26105110

Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Tue, Dec 31, 2019 at 7:38 PM

To: GJESM Journal <gjesm.publication@gmail.com>

Dear Editorial Office,

Thank you for your email.

Here is the authors contributions:

1. Euis Nurul Hidayah: experimental design, analysis, prepare manuscript, literature review
2. Lai Wen Liang: interprete data, prepare manuscript.
3. Okik Hendriyanto Cahyonugroho : arrange data into table and figure, literature review, prepare manuscript
4. Fauzul Rizqa: compiling data and editing manuscript

I would like to ask you to REVISE the name of author : LAI WEN LIANG, which is shown in the article.

LAI is Family Name;

WEN LIANG is his name.

It should be written as W. L. LAI in the article, NOT L.W. LIANG

Thank you

[Quoted text hidden]



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acknowledgement of Revision (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Sat, Nov 30, 2019 at 3:58 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com, nourijafar@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Date: 2019-10-09

Dear **Dr. Euis Nurul Hidayah**

Thank you for submitting the revised file of your manuscript to the **Global Journal of Environmental Science and Management**

The Editorial Office will proceed on your manuscript and inform you in the earliest time.

If there is anything else, please do not hesitate to contact us.

Truly yours,

Professor D. Sivakumar

Managing Editor

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Manuscript Needs Major Revision (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Wed, Nov 27, 2019 at 9:19 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

Your manuscript has now been peer reviewed and virtually resulted as **Major Revision**. As numerous technical comments have pointed out by the reviewers, please revise your manuscript carefully according to the reviewers' comments as well as the editor notifications, within 10 days.

Should the reviewers and editor be satisfied with your amendments, you will be notified the acceptance of your manuscript for publication.

Please do not hesitate to contact us if you have any further inquiries regarding your manuscript revision.

Wishing you every success in your future endeavors.

Truly yours,

Managing Editor

Global Journal of Environmental Science and Management

Dear Author(s),

- 1- Your manuscript is reported as Major revision. The two reviewers reviewed your manuscript and provided scientific comments as attached where can be found through your own dashboard in the website system (#2352).
- 2- Attached, you will also find an "Author Query Form" that you have to add the manuscript reviewer' inquiries and then response each item into the "Author's response" column carefully and correctly in order to be recognized by the reviewers and editor.
- 3- REMEMBER, any alteration and corrections must be done JUST on the attached modified manuscript file (#2352), highlighting with RED fonts in order to be recognized by the Editor and Reviewers.
- 4- After completion of your corrections, please resubmit your revised manuscript as well as the Author Query

Form file through your own dashboard (#2352) where you had already submitted your manuscript in the system as soon as possible to avoid any further delays for the latter processing.

5- Two files are attached in the system as; manuscript #2352 as well as Author Query Form where you have to work on them for your further actions. Every future revisions must be performed on the attached manuscript file not at your previous file.

6- Please also finalize your manuscript English content too.

7- You will have 10 days' time to return your revised manuscript back into correct English. Edition.

Editorial Office

Reviewers Recommendation:

Reviewer 1:

Reviewer Comment For Author:

1. How to determine the characteristic of organic matter? – more and clear information is to be required.
2. During nitrification, the major component of aromatic organic matter changed from low molecular weight of building blocks into high molecular weight of humic substances – justification and explanation are to be required.
3. Batch biofilter nitrification revealed that simultaneous removing the $\text{NH}_4^+\text{-N}$ concentration and increasing organic carbon at the same time revealed microbial products also released by autotrophs during substrate utilization - what type of microbial products generated?
4. What are fractions on obtained components during nitrification process?
5. Comparison of previous studies results with the present study results was missing.
6. There were no distinct results between the two selected methods of study with the combined methods of study - justification is to be required.
7. There was no informative results (values) presented in the abstract – include them.

Reviewer 2:

Reviewer Comment For Author:

Dear Editor

I have read the manuscript carefully. The authors have studied “ Organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix”. As the authors mentioned, lack of studies using both chromatography and fluorescent methods at the same time for characterizing the released organic matter during biological activities, therefore using these two methods to characterize organic matter quantitatively and qualitatively seems to have several potential advantages.

The paper is well written and interesting to read. However, some modifications must be done by the authors to reach a reliable and valuable scientific form of the paper. Herein my main observations:

1. Abstract; do not use abbreviation “FEEM”
2. Abstract; add more data for the results, Abstracts should be able to stand alone.
3. Complete the sentence at the end of introduction
4. Add original references to each section of materials and methods



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Manuscript Needs Resubmission (Manuscript #GJESM-1909-2352 (R1))

4 messages

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Fri, Nov 29, 2019 at 10:20 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com

Dear Author(s),

We are returning your revised paper once more time back! As we have already noticed you, all reviewers' comments and editorial recommendations must be applied at the revised manuscript showing with RED highlighted fonts as some are missing! Also, the performances must be clearly shown and done throughout the text! For example one reviewer had noticed you" In abstract; do not use abbreviation like "FEEM". But you have not followed his comments! and so on. In addition, there are some more remained corrections are shown at the file # 2352 PR3 where all further and final revisions must be done JUST at the mentioned file and returned it back within just 2 more days.

Your delay in resubmitting or uncompleted performances will lead to archiving the manuscript with no further action.

Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Fri, Nov 29, 2019 at 11:57 PM

To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

Dear Editorial Team,

Thank you for your email.

I have gave response to all reviewers' comments in the Author Query Form.

I have checked the file named "2352.doc", only one sentence at page 2 has been highlighted by the red color. I did not find any red highlighted anymore.

I will try to check my account regarding the remained corrections you have mentioned at the new file.

Thank you very much.

=====
Euis Nurul Hidayah, Ph.D.
Assistant Professor
Department of Environmental Engineering
University of Pembangunan Nasional Veteran Jawa Timur
Surabaya, Indonesia
Mobile phone: +6281217870003
=====

[Quoted text hidden]

GJESM Journal <gjesm.publication@gmail.com>

Sat, Nov 30, 2019 at 12:37 AM

To: Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

You have to highlight your corrections throughout the manuscript text body into RED fonts according to the reviewers comments not us!

Truly yours

Editorial Office
Global Journal of Environmental Science and Management (GJESM)

Tel.: +9821- 26105110
Fax: +9821- 26105110

[Quoted text hidden]

Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>
Draft To: GJESM Journal <gjesm.publication@gmail.com>

Sat, Nov 30, 2019 at 12:44 AM

=====
Euis Nurul Hidayah, Ph.D.
Assistant Professor
Department of Environmental Engineering
University of Pembangunan Nasional Veteran Jawa Timur
Surabaya,Indonesia
Mobile phone: +6281217870003
=====

[Quoted text hidden]



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acknowledgement of Revision (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir>

Fri, Nov 29, 2019 at 1:13 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com, nourijafar@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Date: 2019-10-09

Dear **Dr. Euis Nurul Hidayah**

Thank you for submitting the revised file of your manuscript to the **Global Journal of Environmental Science and Management**

The Editorial Office will proceed on your manuscript and inform you in the earliest time.

If there is anything else, please do not hesitate to contact us.

Truly yours,

Professor D. Sivakumar

Managing Editor

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Request for Sending Galley Proof (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir>

Sat, Nov 30, 2019 at 4:56

PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com

(URGENT PLEASE)!

Dear Author,

Attached, you will find three files; 1) Galley Proof, 2) Copyright release, 3) Conflict of interest forms.

Please read the Galley Proof carefully and correct as there are some minor remained revisions which must be corrected. Therefore all new correction must be highlighted in RED fonts (Not yellow paints!) to be recognized by the Editor.

After ending the final correction, complete and sign the two forms of copyrights and return them into PDF format through your dashboard (#2352) in the system immediately for the final processes.

Finally, the 4 Conflict of Interest inquiries must be responded as "No".

Finally, the materials back must be performed by 2 days' time.

Editorial Office



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Submit/Confirm Galley Proof by Author (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir>

Sat, Nov 30, 2019 at 6:57 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

We are pleased to receive your revised Galley Proof along with the signed Copyright release as well as Conflict of Interest forms.

Your manuscript will be processed for publication and the acceptance letter will be forwarded to you shortly.

Truly yours,

Professor J. Nouri

Editor in Cjief

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Acceptance of Manuscript (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Sat, Nov 30, 2019 at 8:02

PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

Acceptance Letter

This is to confirm that after technical and in-house evaluation, the above-mentioned manuscript is finalized and recommended by the Editorial Board Committee to be accepted for publication in the **Global Journal of Environmental Science and Management** (GJESM).

It is necessary to mention that GJESM is a double-blind, peer-reviewed quarterly publication, which is indexed and cited in the well-known world databases mainly at the **Web of Science**, **Scopus**, **SJR (Q2)**, **EBSCO**, **ProQuest**, **Ulrichsweb**, **Cabi**, **Agricola** and **Chemical Abstract**. The title is currently a member of the Committee on Publication Ethics (**COPE**). Upon submission, the manuscript is checked for similarity through a trustworthy software named *iThenticate* to be assured about its originality and then rigorously peer-reviewed by the international reviewers.

Manuscript submission and publication are free of charge in GJESM Journal as a non-commercial publication.

Please remember that upon your published manuscript, it will be accessed and viewed frequently by scholars across the globe. You are therefore recommended to try to enhance its visibility and cite it, not only in your future publications, but also through collaboration with other authors with a similar research theme to promote your research work and attract more citations to your published article(s) in the benefit of your academic career and the GJESM Journal *h-index*. In addition, you may also find your article acceptance certificate at your dashboard.

Thank you for your interest in contributing to the Global Journal of Environmental Science and Management and choosing it as your research hub.

Truly yours,

Professor J. Nouri

Editor in Chief

Global Journal of Environmental Science and Management



Euis Nurul Hidayah <euisnh.tl@upnjatim.ac.id>

Manuscript Published Online (Manuscript #GJESM-1909-2352 (R1))

1 message

Global Journal of Environmental Science and Management <journal@iranjournals.ir> Sat, Nov 30, 2019 at 8:20 PM

Reply-To: Global Journal of Environmental Science and Management <gjesm.publication@gmail.com>

To: euisnh@gmail.com, euisnh.tl@upnjatim.ac.id

Cc: fauzul.rizqa@gmail.com, okikhc.tl@upnjatim.ac.id, laisiff@gmail.com, gjesm.publication@gmail.com, sivakumar.gjesm@gmail.com

Manuscript ID: GJESM-1909-2352 (R1)

Manuscript Title: **Characterization of organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix**

Authors: Euis Nurul Hidayah, Lai Wen Liang, Okik Hendriyanto Cahyo nugroho, Fauzul Rizqa

Dear **Dr. Euis Nurul Hidayah**

I am pleased that your published article is on the net. Kindly visit:

<https://www.gjesm.net/>

Your article is published in the "Article in Press" for the upcoming issues of the Global Journal of Environmental Science and Management (GJESM).

I wish you luck and thank you for sharing your work with us.

Truly yours,

Editorial Office

Global Journal of Environmental Science and Management

Organic matter from biofilter nitrification by high performance size exclusion chromatography and fluorescence excitation-emission matrix

ABSTRACT: [Mw3] [ENH4] A combination of high performance size exclusion chromatography with organic carbon detector and ultraviolet detector coupled with peak-fitting technique and fluorescence [ENH5] excitation-emission matrix spectrometry applied fluorescence regional integration method was conducted to determine the characteristics of organic matter during nitrification. The batch scale of bionet nitrification without organic carbon substrate under aerobic conditions was operated for around 150 minutes. Bulk organic parameters and NH_4^+ -N concentration were analyzed. Five different molecular weights of organic matter were identified by using chromatography, and five different groups of fluorophores organic fractions detected by fluorescence. According to chromatography [ENH6] with carbon and ultraviolet detector, the main characteristics of organic matter shifted from building blocks aromatic compounds with percentage peak area of carbon/ultraviolet detector: 31%/53% to 14%/27.5% to humic-like substances with percentage peak area of carbon/ultraviolet detector 21%/17% to 27%/46.5% during nitrification. Those former compounds are biodegradable [ENH7] as well as properties of microbial products released during substrate utilization and endogenous phase, which are mainly identified as humic-like substances, thus underwent further biodegradation. However, there was significant change in the fluorophores organic fractions, which exhibited humic acid-like with percentage fluorescence regional index area 53% into 68%, as shown by fluorescence excitation-emission matrix analysis. A combination of these methods indicated that the organic matter released during nitrification mainly consists of humic compounds. These results conjecture that a combination of high performance size exclusion chromatography with carbon and ultraviolet detector and fluorescence excitation-emission matrix can be used to determine the characteristic of organic matter and water quality change during nitrification.

KEYWORDS: *Fluorescence excitation-emission matrix (FEEM); Fluorescence regional integration (FRI); Nitrification; Organic matter; Size exclusion chromatography.*

NUMBER OF REFERENCES	NUMBER OF FIGURES	NUMBER OF TABLES
30	5	1

RUNNING TITLE: Characterization organic matter from biofilter

INTRODUCTION

Biological treatment processes, such as bio-filter processes, nitrification, and membrane bioreactor (MBR) release microbial products (Tian *et al.*, 2011; Xie *et al.*, 2016). Although the three types of microbial products, soluble microbial products (SMPs), extracellular polymeric substances (EPS) and inert biomass, are commonly discussed together from a theoretical perspective (Laspidou and Rittman 2002), it is important to distinguish between them because the content of generated organic compound indicated different characteristic then its quality and quality will be vary. The coexistence

of microbial products with natural organic matter (NOM) means that NOM exhibits different qualities during biological treatment, and also increases in quantity (Ni *et al.*, 2011; Liu *et al.*, 2014). After treatment effluent organic matter (EfOM) has been found to contain SMPs. Therefore EfOM in raw water could lead to the formation of disinfection by-products (DBPs) (Liu *et al.*, 2014; Shon *et al.*, 2012). DBPs, which have been found to be carcinogenic and mutagenic, will be formed if EfOM reacts with chlorine or disinfectant during water treatment (Zeng and Mitch 2016). Because the coexistence of heterotrophic bacteria with autotrophs, which increases the organic compound, is potentially problematic, a considerable amount of research has been conducted into the effect of microbial products on the bio-filter nitrification process. Nitrifying bacteria and heterotrophs both need organic substrates to grow, but heterotroph can grow even without a supply of external organic carbon, because they utilize microbial products of nitrifying bacteria, derived from biomass decay and substrate metabolism (Ni *et al.*, 2011; Matsumoto *et al.*, 2010). It is important to understand the characteristics of organic matter in bio-filter nitrification because the variety of organic matter will change due to the effect of microbial products which have been released and/or utilized by the bacteria community in the system, or existing organic carbon (Xie *et al.*, 2016; Ni *et al.*, 2011). Although several studies have investigated the characteristics of microbial products in the form of SMPs and EPS as they undergo biological processes such as activated sludge (Tian *et al.*, 2011; Ni *et al.*, 2011; Kim and Dempsey 2012; Liu *et al.*, 2016; Zhiji *et al.*, 2017), not much work has been conducted in relation to bio-filter nitrification. Organic^[ENH8] matter is complex mixture of heterogeneous compound, which can be classified based on their properties, such as based on molecular weight, aromatic, aliphatic, hydrophilic/hydrophobic, humic/non-humic, etc. Therefore, a number of characterization techniques have been employed to obtain a better understanding of the types of NOM present in source water, and their subsequent removal or transformation through the water treatment process train (Matilainen *et al.*, 2011). High performance size exclusion chromatography (HPSEC) with an on-line organic carbon detector (OCD) and ultraviolet detector (UVD) can detect any type of organic carbon bonded species, this method is based on the molecular size of organic matter. The^[ENH9] separation technique by HPSEC is based on differential permeation of molecules of various size into a porous matrix (Jiao *et al.*, 2014; Lai *et al.*, 2015). Fluorescence spectroscopy, by using fluorescence excitation emission matrices (FEEM), is another option for organic characterization; it can provide insights into the chemical characteristics of NOM because the results are based on both molecular structure and composition (Hidayah *et al.*, 2017). More^[ENH10] rigid aromatic molecular structure and highly conjugated molecules are more likely to fluoresce than aliphatic, alicyclic molecules, and less conjugated systems. Because those molecular structure have smaller energy gaps between the excited and ground states, therefore it will fluoresce at longer wavelength (Murphy *et al.*, 20013). Fluorescence spectroscopy can provide qualitative information to supplement HPSEC, which has limitations as far as the detection of non-chromophores is concerned, and the identification of the chemical and physical properties of the particular molecular size of organic components (Hidayah *et al.*, 2017; Chen *et al.*, 2003). Previous studies have characterized the released organic matter during biological activities by using fluorescent spectroscopy methods only (Liu *et al.*, 2016; Lai *et al.*, 2007; Moradi *et al.*, 2018; Ho *et al.*, 2019; Hidayah and Cahyonugroho 2019). Lack of studies using both chromatography and fluorescent methods at the same time for characterizing the released organic matter during biological activities, therefore using these two methods to characterize organic matter quantitatively and qualitatively seems to have several potential advantages. The objective of this study was to combine the advantages of both the chromatography and the spectrophotometry methods in order to determine the characteristics of organic matter during nitrification and to correlate the effect of ammonia degradation and released organic carbon. In addition, this study was able to quantify the changes in different organic and microbial group compounds in water caused by the nitrification process. This study^[Mw11]^[ENH12] has been carried out in laboratory batch scale at Department of Environmental Engineering, National Cheng Kung University, Taiwan in 2016^[ENH13].

MATERIALS AND METHODS

Bionet filter and sample interval

About 20 pieces small bionet (2 x 4 x 2 cm in size) was put into the sack, it was known as a set of bionet. Bionet is acclimated and taken from nitrification unit in Feng San pilot plant water treatment, then bionet was used as biomedica for nitrifier growth in batch scale process. Bionet is biomass support media with film strips made of UV resistant polyethylene (PEHD) with specific growth surface 100-250 m²/m³. Random media of bionet in has different profiles and large surfaces area allowing the growth and attachment of microorganism. Bionet can be easily removed for replacement or cleaning and bionet sack will not stretch or deteriorate when immersed in water for many years (Jacome *et al.*, 2013). The batch scale of bionet filter made of cylindrical glass was fulfilled with a volume of 2 L synthetic water without organic carbon substrate under aerobic conditions. Bionet was taken from nitrification unit process in Feng-San pilot plant. The batch reactor was operated around 150 minutes in the condition of 30°C. Sampling time was taken from 0 minutes as initial time to interval of 20 minutes for the first hour then interval of 30 minutes for the next hours. Around 20 mL of sample volume were filtered through a 0.45- μ m membrane filter to remove particulate matter.

Synthetic water

Synthetic water contained of ammonia nitrogen of 5 mg-N/L as substrate for autotrophs growth; however, carbon source for heterotrophs growth was not provided in order to characterize released microbial products during nitrification. Bacterial growth required inorganic nutrient and the composition was as follows: K₂HPO₄ 1.3 g/L; KH₂PO₄ 0.3 g/L; MgSO₄·7H₂O 200 mg/L; CaCl₂·2H₂O 20 mg/L; Na₂MoO₄·2H₂O 0.1 mg/L; MnCl₂·4H₂O 0.2 mg/L; ZnSO₄·7H₂O 0.1 mg/L; CuSO₄·5H₂O 0.02 mg/L; CoCl₂·6H₂O 0.002 mg/L; 38.4% Na₂CO₃ 0.1152 g/L; Fe-EDTA solution 1 ml/L (Liu^[ENH14] 2013)

HPSEC, FEEM, NPDOC and NH₄⁺-N analysis

High performance liquid chromatography (HPLC, LC-20 ATV, Shimadzu, Japan) size exclusion chromatography (SEC) was conducted with sequential on-line detectors consisting of a UVD (254 nm, SPD-20A, UV-Vis detector, Shimadzu) and OCD (modified Sievers Total Organic Carbon Analyzer 900 Turbo, General Electric Water & Process Technologies). Fluorescence measurements were undertaken using a Perkin Elmer LS-55 luminescence spectrometer with a xenon lamp and 1.0 cm quartz cell. Excitation emission matrix (EEM) were generated for each sample by scanning over excitation (Ex) wavelengths between 230 and 400 nm at interval of 10 nm and emission (Em) wavelengths between 300 and 547.5 nm at intervals of 0.5 nm (Murphy^[ENH15] *et al.*, 2013; Hidayah *et al.*, 2017). Dissolved organic carbon concentration is represented by non-purgeable dissolved organic carbon (NPDOC) (TOC-5000, Shimadzu, Japan). NH₄⁺-N concentration was measured based on Standard Methods (APHA 2005). Peak fitting technique, PeakFit Version 4.12, Systat Software Inc., USA, was applied to resolve the overlapping peaks of HPSEC chromatogram and to determine the area under each peak. The procedure of peak-fitting technique was described in previous study (Chow^[ENH16] *et al.*, 2008; Lai *et al.*, 2015; Hidayah *et al.*, 2017). Calculation of fluorescence regional integration (FRI) analysis was used with integration beneath EEMs within selected regions to present the cumulative fluorescence response of organic matter with similar properties. The procedure of FRI analysis was described by Chen for analyzing drinking water and wastewater samples on 2003 in Arizona, USA (Chen *et al.*, 2003).

RESULTS AND DISCUSSION

NH₄⁺-N degradation and released organic carbon

As shown in Fig. 1, the nitrification process in bench-scale bionet filter could be divided into two-stage pattern. During the first stage from starting to 40 minutes (min), 90% of 5 mg-NH₃-N/L was utilized by autotrophic microorganisms, indicating that the growth of autotrophic bacteria plays

an important role. At the same stage, the percentage of NPDOC increased to 57.5% from initial value of 2.55 mg-C/L to 6.20 mg-C/L, proving that microbial metabolic products, expressed by NPDOC, was produced in processes of major autotrophs growth or minor heterotrophic metabolism. In addition, carbon source may be related with the diffusion existed in bionet filter taken from Feng-San pilot plant while organic carbon concentration in the bulk phase and was far less than that in bionet filter (Edzwald and Tobiason 2011). Observed from second stage with the slow decomposition of ammonium (40-150 minutes), the accumulation percentage of ammonia removal was close to 99%. The removal ability of autotrophic bacteria on ammonium became slower in comparison with NPDOC removal of 66.4% with extra removal of 8.5%. The released organic carbon was mainly from attributed to microbial products while $\text{NH}_4^+\text{-N}$ substrate utilization was decomposed by autotrophic bacteria. Afterwards, marginal increase of organic carbon was happened while the removal of $\text{NH}_4^+\text{-N}$ slowed down. Ammonia substrate consumption by nitrifier bacteria could lead to microbial products production during in different operational process (Krasner *et al.*, 2013; Xie *et al.*, 2016; Zhiji *et al.*, 2017). Soluble microbial by-products has been released during biological process, including during substrate-utilization associated (microbial growth) and biomass-associated product (during endogenous phase). In general microbial by-product compound is known as humic and fulvic acids, polysaccharides, proteins, nucleic acids, organic acids, amino acids, antibiotics, steroids, exocellular enzymes, siderophores, structural components of cells and products of energy metabolism (Barker and Stuckey 1999). According to Ni *et al.* (2010), it was found that humic-like substances were mainly substrate-utilization associated, while fulvic acid-like substance were non-growth associated, as detected by FEEM. In addition, protein-like substances has been released during microbial growth and polysaccharide-like material has been released during endogenous phase, as detected by Fourier Transform Infra-Red (FTIR). Urbain *et al.* (1998) found that utilization-associate products are mainly carbon compounds generated from the original substrate and that biomass-associated products are cellular macromolecules containing carbon and nitrogen. Briefly, it can be concluded that microbial by-products is composed of different organic compounds.




Fig. 1: The variations of $\text{NH}_4^+\text{-N}$ and NPDOC concentrations during the nitrification operation of bionet filter (Initial ammonium concentration =5 mg/L)

Area percentage of different organic properties from bionet filter

Fig. 2, one of the example HPSEC chromatograms from raw water sample, shows a distribution of organic fractions as detected by HPSEC with OCD and UVD. Peak fitting model of area percentage was applied to chromatogram of each organic fraction from respective measurement by HPSEC with OCD

and UVD during the operation of bionet filter, as shown in Fig. 3. According to the description of previous paragraph, the variation of organic fraction distributions were compared with two-stage growth. Regarding to first stage belonging to rapid decomposition of ammonium or rapid increase of organic carbon, it reveals that the organic fractions were dominated by small molecular weight (MW) of Peak D measured by OCD and UVD, with $250 \text{ Da} < \text{MW} < 500 \text{ Da}$, regarded as low molecular weight acid (Lai *et al.*, 2015; Hidayah *et al.*, 2017). This substance measured by HPSEC-OCD decreased from 31% to 29%, as shown in Fig. 3a, and that measured by UVD descended from 53% to 35.5%, as shown in Fig. 3b. Observed from in Fig. 3a, the increase of biopolymers (Peak A with average molecular weight (AMW) $> 20 \text{ kDa}$) was from 12% to 20% and low molecular weight neutrals (Peak E with AMW $< 250 \text{ Da}$) from 13% to 17.5%. Due to the impurity of EDTA with MW=292, it may be attributed to one of major resource of low molecular weight acid at starting operation. Other organic fractions should be possibly derived from diffusion of organic carbon attached in bionet filter. Release organic matter, including low molecular weight and high molecular weight larger than 10 kDa had been also reported previously (Tian *et al.*, 2011; Laspidou and Rittman 2002; Ni *et al.*, 2011). However, the decrease of building blocks or Peak C from 23% to 14.5% measured by HPSEC-OCD was attributed to the breakdown products from humic substances-like (Huber *et al.*, 2011). As for the second stage, belonging to slow decomposition of ammonium or marginal increase of organic carbon, dominant component was shifted from low molecular weight of acid substance (Peak D) to humic substances-like (Peak B with $500 \text{ Da} < \text{AMW} < 20 \text{ kDa}$). Humic-like substances detected with HPSEC-OCD increased from 18% (40 minutes) to 27% (150 minutes). The similar ascending trend, 29% to 46.5%, was observed by HPSEC-UVD.




Fig. 2: Distribution of organic fractions as detected by HPSEC with (a) OCD and (b) UVD

in water contained of ammonia nitrogen of 5 mg-N/L

The variations of Peak B substance measured by both HPSEC were opposite to those of Peak D. For Peak E substance, the increase of 17.5% to 26 % measured by HPSEC-OCD showed inconsistent with the decrease of HPSEC-UVD from 11% to 3% owing to different organic structure to affect the measurement by HPSEC-OCD and HPSEC-UVD. Several reports demonstrated that HPSEC-OCD could detect organic matter fraction with low absorptivity, such as protein and even fraction with no absorptivity, such as polysaccharide; however, HPSEC-UVD detection is available for the measurement aromatic structure (Huber *et al.*, 2011; Lai *et al.*, 2015; Hidayah *et al.*, 2017). The close increase of 20% to 21% for Peak A by OCD in relative with UVD showed same pattern with Peak E, indicating that the organic structures of Peak E and Peak A contained less aromatic or unsaturated function group. Moradi^[ENH18] *et al.* (2017) used HPSEC with UVD at wavelength 230 nm to investigate the nitrification occurrence in two drinking water distribution systems in Australia, and the results shows that formation of soluble microbial products and/or the release of extracellular polymeric substances (EPS) during nitrification. A peak of EPS has appeared at apparent molecular weight < 500 Da, which is similar peak at this present study. The peak less than 500 Da are identified as humic substances-like (Peak B) and building block (Peak C) in this present study. Regarding to the processes of nitrification, the transformation of organic compounds fraction was significantly affected by nitrifer to utilize ammonia (Xie *et al.*, 2016; Ni *et al.*, 2011; Matsumoto *et al.*, 2010). Briefly^[ENH19], HPSEC with OCD and UVD has been identified 5 fractions of organic matter in water sampel and during nitrification. It can concluded that humic substances-like has been generated during nitrification process, as the initial stage showed 21% area of OCD and 17% area of UVD become increased into 27% area of OCD and 46.5% area of UVD.




Fig. 3: Area percentage of organic fraction distribution during the bionet expressed by (a) HPSEC-OCD and (b) HPSEC-UVD with peakfitting software

Variations of organic FEEM

Fig. 4, one of the example FEEM spectra from raw water sample, shows a distribution of organic fractions as detected by fluorescence excitation-emission matrix (FEEM) spectroscopy. FEEM was applied to characterize the properties of organic matter through bionet system. Five organics classification were divided from the FEEM contour according to the previous research (Chen *et al.*, 2003). Region I, located at shorter excitation/emission wavelengths (Ex/Em nm), <250/<330 nm, is known as simple aromatic proteins such as tyrosine. Region II, located at <250/330-380 nm, is related to simple aromatic protein such as tryptophan. Region III, located at 200-250/>380 nm, is

identified as fulvic like substances. Excitation/emission wavelengths of 250-280/<380 nm related to soluble microbial by product-like material is classified as Region IV. Peak at longer excitation/emission wavelengths of >280/>380 nm, named as Region V, is represented to humic acid-like substances. Average fluorescent intensities of five regions of organic fractions were calculated by FRI methods according to the method proposed in previous study (Chen *et al.*, 2003; Guo *et al.*, 2015).




Fig. 4: Distribution of organic fractions as detected by fluorescence excitation-emission matrix (FEEM) spectroscopy

As shown in Fig. 5, the FEEM classification of organic matter from during nitrification was happened in batch-scale bionet filter. For the starting operation, 53% of humic acid-like proved a fact that bionet filter taken from on-site biofilter contains the accumulated refractory organic matter. Of course, EDTA with aminocarboxylic acid, belonging to humic like polycarboxylate-type, was one of contributions (Liu *et al.*, 2014; Liu *et al.*, 2016) however, the dominant fractions may be related with the diffusion from exhausted bionet filter (Han *et al.*, 2013; Greenstein *et al.*, 2018). After the operation of 40 minutes, fluorescent characteristic released from bionet filter was mainly occupied by humic like substance of 65%, soluble microbial-products like tryptophan of 12%, fulvic-like substance of 18% and aromatic protein of 7%. This FRI percentage of different organic property was kept constant at 150 minutes of operation. Compared starting operation with 40 minutes, the decrease of fulvic-like substance was opposite to the increase of humic-like substance, revealing that humic-like substance is one of accumulated material while biodegradable process is operated in the bionet filter. Similar reports were demonstrated during the substrate consumption by microorganisms (Ni *et al.*, 2011; Liu *et al.*, 2014; Liu *et al.*, 2016). Briefly (ENH20), FEEM with FRI analysis has been identified 5 fractions of organic matter in water sample and during nitrification. It can concluded that humic acid-like has been generated during nitrification process, as the initial stage showed 53% area of FRI increased become 68% area of FRI. The results is consistent with increasing of humic substances-like as detected by HPSEC-OCD and-UVD with peak-fitting technique.




Fig. 5: Variations of average FRI values and percentages for different organic fraction distribution during bionet operation

Accumulation of humic-like substance as long-term operation of bionet

According to the current results, the increase of organic carbon was not only related with diffusion of organic originated from bionet media but also with metabolic matter from heterotrophic bacteria located outside of the bionet media while bionet filter was initially operated. In the nitrification process, autotrophic bacteria had gradually grown and eventually present in the outer part of the bionet media (Matsumoto *et al.*, 2010; Chang *et al.*, 2019), implying that in this study, fast and slow ammonium decomposition may be attributed to the competition between ammonium nitrifying bacteria and heterotrophic bacteria. Furthermore, released organic fractions could be affected by substrate utilization, transformation to cell mass and metabolic product by autotrophic and heterotrophic microorganism (Laspidou and Rittman 2002; Ni *et al.*, 2011). Previous studies (Liu *et al.*, 2014; Lai *et al.*, 2007; Moradi *et al.*, 2018) also used FEEM to characterize organic fractions and found protein, polyaromatic humic acid-like and polycarboxylate humic acid-like substances were produced and accumulated while the operation time was continuously conducted in the biofilter. Those previous studies had identified humic-acid-like organics as shown a significant peak at Ex/Em wavelength 330/435 nm (Liu *et al.*, 2014), 325/420 nm (Lai *et al.*, 2007), 325/425 nm (Moradi *et al.*, 2018), which are similar peak with this present study. In slow ammonium removal, increased NPDOC removal was higher than ammonium removal. The microbial products released in the later phase of biodegradation has high molecular weight because the substrates have been exhausted or depleted, resulting that microorganisms were in decay phase (Ni *et al.*, 2011; Liu *et al.*, 2016). Low molecular weight acid (Peak D) and building blocks (Peak C) decrease during slow ammonium decomposition because those compounds are biodegradable as well as properties of microbial products released during substrate utilization and thus underwent further biodegradation. Heterotrophs could utilized and degraded some released microbial by-product including low molecular weight acid component (Ni *et al.*, 2011; Matsumoto *et al.*, 2010), since heterotrophs existed in bionet filter and survived by background organic carbon from preparation of water sample and diffusion from bionet filter. According to Ni *et al.* (2010), it was found that humic-like substances were mainly substrate-utilization associated, while fulvic acid-like substance were non-growth associated, and both fractions is characterized as humic substances-like.

Table 1: Variation of five organic fractions expressed by HPSEC-OCD, HPSEC-UVD and FEEM from bionet filter operation with nitrification processes

Methods	Operation time (minutes)	Increase (%) [*]					Total
		Peak A	Peak B	Peak C	Peak D	Peak E	
HPSEC-OCD	0 to 40	14	7	2.2	12.5	10.5	46.2
	40 to 150	17.3	21.5	4.5	3.4	20.5	67.2
HPSEC-UVD	0 to 40	0.3	19	8.4	3.3	8.3	39.3
	40 to 150	1.4	38.1	8.8	1.5	1.1	50.9
FEEM		Region 1	Region 2	Region 3	Region 4	Region 5	
	0 to 40	0.2	1	9.3	2.6	41	54.1
	40 to 150	0.3	1.6	10.1	3.2	49	64.2

^{*} For HPSEC = [(Peak Area_{operation time} – Peak Area_{starting})/Peak Area_{operation time}] x 100%

For FEEM = [(Region_{operation time} – Region_{starting})/Region_{operation time}] x 100%

Table 1 lists the variations of released organic properties from bionet filter with nitrification expressed by HPSEC-OCD, HPSEC-UVD and FEEM. At fast ammonium decomposition, HPSEC-OCD showed that biopolymers (i.e., protein) or Peak A, low molecular weight acids or Peak D, and Peak E of low molecular weight neutrals respectively had the increase of 14%, 12.5% and 10.5%, indicating that the dominant substances were Peak A, D and E, however, the increase of 19% for humic substances-like became more prominent by the measurement of HPSEC-UVD. This discrepancy was due to UV sensible for the detection of aromatic unsaturated groups (Lai *et al.*, 2015; Hidayah *et al.*, 2017) in opposite to difficult combustion by OCD. Observed from FEEM during fast ammonium decomposition, the increase of humic like substance was 41% higher than other organic fraction (i.e., protein and soluble microbial products as microbial by-product). Overall, the increase of whole organic compounds during 40 minutes was in the order of 54% for FEEM, 46% from HPSEC-OCD, and 39% for HPSEC-UVD. Regarding the characteristic of released organic matter at slow ammonium decomposition, Peak B owned more important role than Peak A and E whatever HPSEC-OCD or HPSEC-UVD was applied. Peak B as microbial by-products is hard to be degraded and will be accumulated in the system. Overall, the whole organic compounds during 110 minutes was in the order, 67% of HPSEC-OCD, 51% of HPSEC-UVD, and 64% of FEEM. Obviously, organic fractions increased marginally around 19% - 39% by HPSEC and 23% by FEEM during 110 minutes. Listed in Table 1, during substrate depletion, polycarboxylate humic acid-like and protein kept increasing trend with consistence (Li *et al.*, 2013; Liu *et al.*, 2014). Those humic-like is non-biodegradable compounds (Ni *et al.*, 2011; Zhiji *et al.*, 2017), therefore, being a dominant component in the second stage is reasonable.

FEEM and HPSEC with both detectors have characterized that all organic fractions increased and identified the same fractions during fast and slow ammonium decomposition. In this study, combining the derived fraction obtained from HPSEC-OCD and -UVD with peak-fitting techniques and the derived component from F-EEM with FRI analysis, on the same sample simultaneously, gave a reinforced information and had confirmed the presence of humic substances-like (Peak B)/humic acid-like (Region 5) as the main organic in nitrification processes. The study indicates that a combination of HPSEC-OCD/UVD and FEEM can be used to determine the characteristics of organic matter and water quality change during nitrification, though FEEM seemed like having more sensitivity than the other instruments.

CONCLUSION

Application of HPSEC-OCD/UVD with peak-fitting method and FEEM with FRI method identified the characteristics of organic compounds released during nitrification. HPSEC-OCD classified five different types of organic matter based on average molecular weight, including biopolymers, humic substances,

building blocks, low molecular weight acid, and low molecular weight neutral. HPSEC-UVD showed that these organic compounds have an aromatic structure. FEEM with FRI technique characterized five different organic fluorophores: aromatic protein 1, aromatic protein 2, fulvic acid-like, humic acid-like, and soluble microbial products. During nitrification, the major component of aromatic organic matter changed from low molecular weight of building blocks into high molecular weight of humic substances. This is consistent with the dominant compounds humic acid-like substances in FEEM analysis. In addition, batch biofilter nitrification revealed that simultaneous removing the $\text{NH}_4^+\text{-N}$ concentration and increasing organic carbon at the same time revealed microbial products also released by autotrophs during substrate utilization or in the stage of ammonium decomposition. These results lead to the conjecture that the organic matter released during nitrification is mainly characterized by humic substances compounds. The study indicates that a combination of HPSEC-OCD/UVD and FEEM can be used to determine the characteristics of organic matter and water quality change during nitrification.

ACKNOWLEDGEMENT

Authors would like to thank to Environmental Biotechnology Laboratory, Department of Environmental Engineering, National Cheng Kung University for supporting experiment.

CONFLICT OF INTEREST

The author declares that there is no conflict of interests regarding the publication of this manuscript. In addition, the ethical issues, including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy have been completely observed by the authors.

ABBREVIATIONS^[Mw24]^[ENH25]

%	Percentage
μm	Micrometer
AMW	Average molecular weight
C	Carbon
cm	centimeter
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	Calcium chloride dihydrate
$\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$	Cobaltous chloride hexahydrate
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	Copper sulfate pentahydrate
Da	Dalton
DBPs	Disinfection by-products
EDTA	Ethylene diamine tetra acetic acid
EEM	Excitation-emission matrix
EfOM	Effluent organic matter
Em	Emission
EPS	Extracellular polymeric substances
Ex	Excitation
Fe	Ferric

<i>Fe-EDTA</i>	Ferric ethylene diamine tetra acetic acid
<i>FEEM</i>	Fluorescence excitation-emission matrix
<i>FRI</i>	Fluorescence regional integration
<i>FTIR</i>	Fourier transform infra red
<i>g/L</i>	Gram per-liter
<i>H₂O</i>	Hydrogen dioxygen
<i>HPSEC</i>	High performance size exclusion chromatography
<i>HPSEC-OCD</i>	High performance size exclusion chromatography organic carbon detector
<i>HPSEC-UVD</i>	High performance size exclusion chromatography ultra violet detector
<i>K₂HPO₄</i>	Dipotassium phosphate
<i>KH₂PO₄</i>	Potassium dihydrogen phosphate
<i>kDa</i>	Kilo dalton
<i>L</i>	Liter
<i>m²/m³</i>	Square meter per-cubic meter
<i>MBR</i>	Membrane bioreactor
<i>MgSO₄ · 7H₂O</i>	Magnesium sulfate heptahydrate
<i>mg/L</i>	Milligram per-liter
<i>mg-C/L</i>	Miligram carbon per-liter
<i>mg-NH₃-N/L</i>	Miligram ammonia nitrogen per-liter
<i>min</i>	Minutes
<i>mL</i>	Milliliter
<i>ml/L</i>	Milliliter per-liter
<i>MnCl₂ · 4H₂O</i>	Manganese (II) chloride tetrahydrate
<i>MW</i>	Molecular weight
<i>Na₂CO₃</i>	Sodium carbonate
<i>Na₂MoO₄ · 2H₂O</i>	Sodium molybdate dihydrate
<i>NH₃-N</i>	Ammonia-nitrogen
<i>NH₄⁺-N</i>	Ammonium-nitrogen
<i>nm</i>	Nanometer
<i>NOM</i>	Natural organic matter
<i>NPDOC</i>	Non-purgeable dissolved organic carbon
<i>PEHD</i>	Polyethylene high density
<i>OCD</i>	Organic carbon detector
<i>SMPs</i>	Soluble microbial products
<i>TOC</i>	Total organic carbon
<i>USA</i>	United states of america
<i>UV</i>	Ultraviolet
<i>UV-Vis</i>	Ultra violet visible

UVD	Ultraviolet detector
ZnSO ₄ · 7H ₂ O	Zinc sulfate heptahydrate

REFERENCES

APHA, (2005). Standard methods for the examination of water and wastewaters, 21th edition, APHA, AWWA, WEF Publication, Washington D.C.

[https://www.scirp.org/\(S\(czeh2tfqyw2orz553k1w0r45\)\)/reference/ReferencesPapers.aspx?ReferenceID=1870039](https://www.scirp.org/(S(czeh2tfqyw2orz553k1w0r45))/reference/ReferencesPapers.aspx?ReferenceID=1870039)

Barker, D.J.; Tuckey, D.C., (1999). A review of soluble microbial products (smp) in wastewater treatment systems. *Water Res.*, 30(14): 3063-3082 (20 pages).

<https://www.sciencedirect.com/science/article/abs/pii/S0043135499000226>

Chang, J.J.; Mei, J.; Jia, W.; Chen, J.Q.; Li, X.; Ji, B.H.; Wu, H.M., (2019). Treatment of heavily polluted river water by tidal-operated biofilters with organic/inorganic media: evaluation of performance and bacterial community. *Biores. Technol.*, 279: 34-42 (9 pages)

<https://www.sciencedirect.com/science/article/pii/S0960852419300732>

Chen, W.; Westerhoff, P.; Leenheer, J.A.; Booksh, K., (2003). Fluorescence excitation-emission matrix regional integration to quantify spectra for dissolved organic matter. *Environ. Sci. Technol.*, 37(24): 5701-5710 (10 pages).

<https://asu.pure.elsevier.com/en/publications/fluorescence-excitation-emission-matrix-regional-integration-to-q>

Chow, C. W. K.; Fabris, R.; van Leeuwen, J.; Wang, D. S.; Drikas, M., (2008). Assessing natural organic matter treatability using high performance size exclusion chromatography. *Environ. Sci. Technol.*, 42: 6683-6689 (7 pages).

<https://pubs.acs.org/doi/abs/10.1021/es800794r>

Edzwald, J.K.; Tobiason, J.E., (2011). Chemical principles, source water composition, and watershed protection. *Water quality and treatment: a handbook on drinking water*, AWWA, McGraw-Hill, New York.

<https://www.accessengineeringlibrary.com/content/book/9780071630115/chapter/chapter3>

Greenstein, K.E.; Lew, J.; Dickenson, E.R.V.; Wert, E.C., (2018). Investigation of biotransformation, sorption, and desorption of multiple chemical contaminants in pilot-scale drinking water biofilters. *Chemosphere*. 200: 248-256 (9 pages)

<https://www.sciencedirect.com/science/article/pii/S0045653518303199>

Guo, L.; Lu, M.M.; Li, Q.Q.; Zhang, J.W.; She, Z.L., (2015). A Comparison of different pretreatments on hydrogen fermentation from waste sludge by fluorescence excitation-emission matrix with regional integration analysis. *Int. J. Hydrogen Energy*. 40 (1): 197-208 (12 pages)

<https://www.sciencedirect.com/science/article/abs/pii/S0360319914030171>

Han, Z.Y.; Liu, D.; Li, Q.B., (2013). A removal mechanism for organics and nitrogen in treating leachate using a semi-aerobic aged refuse biofilter. *J. Environ. Manage.*, 114: 336-342 (7 pages)

<https://www.sciencedirect.com/science/article/pii/S0301479712005300>

Hidayah, E.N.; Chen, Y.C.; Yeh, H.H., (2017). Comparison between HPSEC-OCD and F-EEMs for assessing DBPs formation in water. *J. Environ. Sci. Health. Part A.*, 52(4): 391-402 (12 pages).

<https://www.tandfonline.com/doi/abs/10.1080/10934529.2016.1262607?scroll=top&needAccess=true&journalCode=lesa20>

Hidayah, E.N.; Cahyonugroho, O.H., (2019). Tracking of dissolved effluent organic matter (dEfOM) in wastewater treatment plant by using fluorescence method. IOP Conference Series: Earth and Environmental Science, 245(1).

<https://iopscience.iop.org/article/10.1088/1755-1315/245/1/012018>

Ho, H.J.; Cao, J.W.; Kao, C.M.; Lai, W.L., (2019). Characterization of released metabolic organic during AOC analyses by P17 and NOX strains using 3-D fluorescent signals. Chemosphere. 222: 205-213 (9 pages).

<https://www.sciencedirect.com/science/article/pii/S0045653519301201>

Huber, S.A.; Balz, A.; Abert, M.; Pronk, W., (2011). Characterisation of aquatic humic and non-humic matter with size-exclusion chromatography-organic carbon detection- organic nitrogen detection (LC-OCD-OND). Water Res., 45: 879-885 (7 pages).

<https://www.sciencedirect.com/science/article/abs/pii/S0043135410006779>

Jacome, A.; Molina, J.; Novoa, R.; Suarez, J.; Ferreira, S., (2013). Simultaneous carbon and nitrogen removal from municipal wastewater in full-scale unaerated/aerated submerged filters. Water Sci. Technol., 69 (1): 217-221 (5 pages)

<https://iwaponline.com/wst/article/69/1/217/18183/Simultaneous-carbon-and-nitrogen-removal-from>

Jiao, R.Y.; Chow, C.W.K.; Xu, H.; Yang, X.F.; Wang, D.S., (2014). Organic removal assessment at full-scale treatment facilities using advanced organic characterization tools. Environ. Sci. Processes Impacts, 16: 2451-2459 (9 pages)

<https://pubs.rsc.org/en/content/articlehtml/2014/em/c4em00227j>

Kim, H.C.; Dempsey, B.A., (2012). Comparison of two fractionation strategies for characterization of wastewater effluent organic matter and diagnosis of membrane fouling. Water Res., 46: 3714-3722 (9 pages).

<https://www.sciencedirect.com/science/article/abs/pii/S0043135412002801>

Krasner, S.W.; Mitch, W.A.; McCurry, D.L.; Hanigan, D.; Westerhoff, P., (2013). Formation, precursors, control, and occurrence of nitrosamines in drinking water: A review. Water Res., 47 (13): 4433-4450 (18 pages).

<https://www.sciencedirect.com/science/article/abs/pii/S004313541300393X>

Lai, W.L.; Chen, L.F.; Liao, S.W.; Hsu, S.L.; Tseng, L.H.; Miaw, C.L., (2007). Using EEFM (excitation emission fluorescence matrix) to differentiate the organic properties of the effluents from the ozonated biofilters. Water Air Soil. Pollut., 186(1-4): 43-53 (11 pages).

<https://link.springer.com/article/10.1007/s11270-007-9461-6>

Lai, C.H.; Chou, Y.C.; Yeh, H.H., (2015). Assessing the interaction effects of coagulation pretreatment and membrane material on UF fouling control using HPSEC combined with peak-fitting. J. Membrane Sci., 474: 207-214 (8 pages).

<https://www.sciencedirect.com/science/article/pii/S0376738814007546>

Laspidou, C.S.; Rittmann, B.E., (2002). A unified theory for extracellular polymeric substances, soluble microbial products, and active and inert biomass. Water Res., 36: 2711-2720 (10 pages).

<https://www.sciencedirect.com/science/article/abs/pii/S0043135401004134>

Li, W.T.; Xu, Z.X.; Li, A.M.; Wu, W.; Zhou, Q.; Wang, J.N., (2013). HPLC/HPSEC-FLD with multi-excitation/emission scan for EEM interpretation and dissolved organic matter analysis. *Water Res.*, 47 93): 1246-1256 (**11 pages**)

<https://www.sciencedirect.com/science/article/abs/pii/S0043135412008524>

Liu,^[Mw26]^[ENH27] Y.W., (2013). Pilot-study on nitrification performance of bionet for ammonia removal in raw water of drinking water treatment. Master Thesis. Department of Environmental Engineering, National Cheng University, Taiwan (**105 pages**).

http://etds.lib.ncku.edu.tw/etdservice/view_metadata?etdun=U0026-3107201415043300

Liu, J.L.; Li, X.Y.; Xie, Y.F.; Tang, H., (2014). Characterization of soluble microbial products as precursors of disinfection byproducts in drinking water supply. *Sci. Total Environ.*, 472: 8818-8824 (**7 pages**).

<https://www.sciencedirect.com/science/article/pii/S0048969713014435>

Liu, Y.; Sun, J.; Peng, L.; Wang, D.; Dai, X.; Ni, B.J., (2016). Assessment of heterotrophic growth supported by soluble microbial products in anammox biofilm using multidimensional modeling. *Sci Rep.*, 6:27576 (**11 pages**).

<https://www.nature.com/articles/srep27576>

Matilainen, A.; Gjessing, E.T.; Lahtinen, T.; Hed, L.; Bhatnagar, A.; Sillanpaa, M., (2011). An overview of the methods used in the characterization of natural organic matter (NOM) in relation to drinking water treatment. *Chemosphere*. 83 (11): 1431-1442 (**12 pages**)

<https://www.sciencedirect.com/science/article/pii/S0045653511000208>

Matsumoto, S.; Kotaku, M.; Saeki, G.; Terada, A.; Aoi, Y.; Tsuneda, S.; Picioreanu, C.; van Loosdrecht, M.C.M., (2010). Microbial community structure in autotrophic nitrifying granules characterized by experimental and simulation analyses. *Environ. Microbiol.* 12(1):192-206 (**15 pages**).

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1462-2920.2009.02060.x>

Moradi, S.; Liu, S.; Chow, C.W.K.; van Leeuwen, J.; Cook, D.; Drikas, M.; Amal, R., (2017). Developing a chloramine decay index to understand nitrification: a case study of two chloraminated drinking water distribution systems. *J. Environ. Sci.*, 57: 170-179 (**10 pages**).

<https://www.sciencedirect.com/science/article/pii/S1001074216312359>

Moradi, S.; Sawade, E.; Aryal, R.; Chow, C.W.K.; van Leeuwen, J.; Drikas, M.; Cook, D.; Amal, R., (2018). Tracking changes in organic matter during nitrification using fluorescence excitation-emission matrix spectroscopy coupled with parallel factor analysis (feem/parafac). *J. Environ. Chem. Eng.*, 6(1): 1522-1528 (**7 pages**).

<https://www.sciencedirect.com/science/article/pii/S2213343718300757>

Murphy, K.R.; Stedmon, C.A.; Graeber, D.; Bro, R., (2013). Fluorescence spectroscopy and multi-way techniques. *PARAFAC. Anal. Methods*, 5(23): 6557-6566 (**10 pages**).

<https://pubs.rsc.org/en/content/articlelanding/2013/ay/c3ay41160e#!divAbstract>

Ni, B.J.; Zeng, R.J.; Fang, F.; Xie, W.M.; Sheng, G.P.; Yu, H.Q., (2010). Fractionating soluble microbial products in the activated sludge process. *Water Res.*, 44: 2292-2302 (**11 pages**).

<https://www.sciencedirect.com/science/article/abs/pii/S0043135409008379>


Ni, B.J.; Xie, W.M.; Chen, Y.P.; Fang, F.; Liu, S.Y.; Ren, T.T.; Sheng, G.P.; Yu, H.Q.; Liu, G.; Tian, Y.C., (2011).

- Heterotrophs grown on the soluble microbial products (SMP) released by autotrophs are responsible for the nitrogen loss in nitrifying granular sludge. *Biotechnol. Bioeng.*, 108(12): 2844-2852 (**9 pages**).
<https://onlinelibrary.wiley.com/doi/full/10.1002/bit.23247>
- Shon, H.K.; Vigneswaran, S.; Snyder, S.A., (2012). Effluent organic matter (EfOM) in wastewater: constituents, effect, and treatment. *Crit. Rev. Environ. Sci. Technol.*, 36(4): 327-374 (**48 pages**).
<https://www.tandfonline.com/doi/full/10.1080/10643380600580011>
- Tian, Y.; Chen, L.; Jiang, T.L., (2011). Characterization and modeling of the soluble microbial products in membrane bioreactor. *Sep. Purif. Technol.*, 70: 316-324 (**9 pages**).
<https://www.sciencedirect.com/science/article/pii/S1383586610004454>
- Urbain, V.; Mobarry, B.; de Silva, V.; Stahl, D. A.; Rittmann, B. E.; Manem, J., (1998). Integration of performance, molecular biology and modeling to describe the activated sludge process. *Water Sci. Technol.*, 37: 223-229 (**7 pages**).
<https://www.sciencedirect.com/science/article/abs/pii/S0273122398001115>
- Zeng, T.; Mitch, W.A., (2016). Impact of nitrification on the formation of N-nitrosamines and halogenated disinfection byproducts within distribution system storage facilities. *Environ. Sci. Technol.*, 50 (6): 2964-2973 (**10 pages**)
<https://pubs.acs.org/doi/10.1021/acs.est.5b05668>
- Xie, W.M.; Ni, B.J.; Sheng, G.P.; Seviour, T.; Yu, H.Q., (2016). Quantification and kinetic characterization of soluble microbial products from municipal wastewater treatment plants. *Water Res.*, 88: 703-710 (**8 pages**).
<https://www.sciencedirect.com/science/article/abs/pii/S0043135415303274>
- Zhiji, D.; Bourven, I.; van Hullebusch; Panico, A.; Pirozzi, F.; Esposito, G.; Guibaud, G., (2017). Quantitative and qualitative characterization of extracellular polymeric substances from Anammox enrichment. *J. Taiwan Inst. Chem. Eng.*, 80: 738-746 (**9 pages**).
<https://www.sciencedirect.com/science/article/pii/S1876107017304765>

HIGHLIGHTS

- Combination of HPSEC-OCD/UVD and FEEM can be used to determine the characteristic of organic matter and water quality change during nitrification;
- During nitrification, the major component of aromatic organic matter changed from low molecular weight of building blocks into high molecular weight of humic substances;
- Simultaneous removing the $\text{NH}_4^+\text{-N}$ concentration and increasing organic carbon at the same time revealed microbial products also released, mainly humic substances fraction.

GRAPHICAL ABSTRACT



Manuscript # 2352

Author Query Form

Add the manuscript reviewing comments of each reviewers into the “details reviewing required” separately and then response each items into the “Author’s response” columns in order to be recognized by the reviewers and editor.

Reviewer # 1:

Query	Details reviewing required	Author’s response (along with the place of corrections in manuscript)
1.	How to determine the characteristic of organic matter? – more and clear information is to be required.	<p>A number of characteristic of organic matter have been added in the revised manuscript. Please see:</p> <p>Abstract Line 1-Line 4</p> <p>A combination of high performance size exclusion chromatography with organic carbon detector and ultraviolet detector coupled with peak-fitting technique and fluorescence excitation-emission matrix spectrometry applied fluorescence regional integration method was conducted to determine the characteristics of organic matter during nitrification.</p> <p>Please see the red highlighted of Introduction Page 2, Line 18-Line 39.</p> <p>Organic matter is complex mixture of heterogeneous compound, which can be classified based on their properties, such as based on molecular weight, aromatic, aliphatic, hydrophilic/hydrophobic, humic/non-humic, etc. Therefore, a number of characterization techniques have been employed to obtain a better understanding of the types of NOM present in source water, and their subsequent removal or transformation through the water treatment process train (Matilainen <i>et al.</i>, 2011). High performance size exclusion chromatography (HPSEC) with an on-line organic carbon detector (OCD) and ultraviolet detector (UVD) can detect any type of organic carbon bonded species, this method is based on the molecular size of organic matter. The separation technique by HPSEC is based on differential permeation of molecules of various size into a porous matrix (Jiao <i>et al.</i>, 2014; Lai <i>et al.</i>, 2015). Fluorescence spectroscopy, by using fluorescence excitation emission matrices (FEEM), is another option for organic characterization; it can provide insights into the chemical characteristics of NOM because the results are based on both molecular structure and composition (Hidayah <i>et al.</i>, 2017). More rigid aromatic molecular structure and highly conjugated molecules are more likely to fluoresce than aliphatic, alicyclic molecules, and less conjugated systems. Because those molecular structure have smaller energy gaps between the excited and ground states, therefore it will fluoresce at longer wavelength (Murphy <i>et al.</i>, 20013). Fluorescence spectroscopy can provide qualitative information to supplement HPSEC, which has limitations as far as the detection of non-chromophores is concerned, and the identification of the chemical and physical properties of the particular molecular size of organic components (Hidayah <i>et al.</i>, 2017; Chen <i>et al.</i>, 2003).</p>

<p>2.</p>	<p>During nitrification, the major component of aromatic organic matter changed from low molecular weight of building blocks into high molecular weight of humic substances – justification and explanation are to be required.</p>	<p>Explanation has been added in the revised manuscript.</p> <p>Please see the red highlighted of Abstract Line 9-Line 15. According to chromatography with carbon and ultraviolet detector, the main characteristics of organic matter shifted from building blocks aromatic compounds with percentage peak area of carbon/ultraviolet detector: 31%/53% to 14%/27.5% to humic-like substances with percentage peak area of carbon/ultraviolet detector 21%/17% to 27%/46.5% during nitrification. Those former compounds are biodegradable as well as properties of microbial products released during substrate utilization and endogenous phase, which are mainly identified as humic-like substances, thus underwent further biodegradation.</p> <p>Detail explanation has been mentioned in the manuscript Result and Discussion, Page 8, Line 18 – Line 24 and additional explanation with the red highlighted Page 8 Line 25 – Line 27. Low molecular weight acid (Peak D) and building blocks (Peak C) decrease during slow ammonium decomposition because those compounds are biodegradable as well as properties of microbial products released during substrate utilization and thus underwent further biodegradation. Heterotrophs could utilized and degraded some released microbial by-product including low molecular weight acid component (Ni <i>et al.</i>, 2011; Matsumoto <i>et al.</i>, 2010), since heterotrophs existed in bionet filter and survived by background organic carbon from preparation of water sample and diffusion from bionet filter. According to Ni <i>et al.</i> (2010), it was found that humic-like substances were mainly substrate-utilization associated, while fulvic acid-like substance were non-growth associated, and both fractions is characterized as humic substances-like.</p>
<p>3.</p>	<p>Batch biofilter nitrification revealed that simultaneous removing the NH₄⁺-N concentration and increasing organic carbon at the same time revealed microbial products also released by autotrophs during substrate utilization - what type of microbial products generated?</p>	<p>Type of microbial by-products has been added in the manuscript. Please see the red highlighted of: Results and discussion Page 4, line 11– line 23 Soluble microbial by-products has been released during biological process, including during substrate-utilization associated (microbial growth) and biomass-associated product (during endogenous phase). In general microbial by-product compound is known as humic and fulvic acids, polysaccharides, proteins, nucleic acids, organic acids, amino acids, antibiotics, steroids, exocellular enzymes, siderophores, structural components of cells and products of energy metabolism (Barker and Stuckey 1999). According to Ni <i>et al.</i> (2010), it was found that humic-like substances were mainly substrate-utilization associated, while fulvic acid-like substance were non-growth associated, as detected by FEEM. In addition, protein-like substances has been released during microbial growth and polysaccharide-like material has been released during endogenous phase, as detected by FTIR. Urbain <i>et al.</i> (1998) stated that utilization-associate products are mainly small carbonaceous compounds derived from the original substrate and that biomass-associated products are cellular macromolecules containing both carbon and nitrogen. Briefly, it can be concluded that microbial by-products is composed of different organic compounds.</p>

4.	What are fractions on obtained components during nitrification process?	<p>Obtained fractions during nitrification has been concluded in the manuscript. Please see the red highlighted of:</p> <p>Results and discussion Page 6, Line 17 – Line 21 Briefly, HPSEC with OCD and UVD has been identified 5 fractions of organic matter in water sampel and during nitrification. It can concluded that humic substances-like has been generated during nitrification process, as the initial stage showed 21% area of OCD and 17% area of UVD become increased into 27% area of OCD and 46.5% area of UVD.</p> <p>Results and discussion Page 7, Line 20 – Line 24 Briefly, FEEM with FRI analysis has been identified 5 fractions of organic matter in water sampel and during nitrification. It can concluded that humic acid-like has been generated during nitrification process, as the initial stage showed 53% area of FRI increased become 68% area of FRI. The results is consistent with increasing of humic substances-like as detected by HPSEC-OCD and-UVD with peak-fitting technique.</p>
5.	Comparison of previous studies results with the present study results was missing.	<p>Comparison of previous studies results with present study has been added in the manuscript. Please see the red highlighted of:</p> <p>Results and discussion Page 6, Line 10- Line 15. Moradi <i>et al.</i> (2017) used HPSEC with UVD at wavelength 230 nm to investigate the nitrification occurrence in two drinking water distribution systems in Australia, and the results shows that formation of soluble microbial products and/or the release of extracellular polymeric substances (EPS) during nitrification. A peak of EPS has appeared at apparent molecular weight < 500 Da, which is similar peak at this present study. The peak less than 500 Da are identified as humic substances-like (Peak B) and building block (Peak C) in this present study.</p> <p>Results and discussion Page 8, Line 14 – Line 16 Those previous studies had identified humic-acid-like organics as shown a significant peak at Ex/Em wavelength 330/435 nm (Liu <i>et al.</i>, 2014), 325/420 nm (Lai <i>et al.</i>, 2007), 325/425 nm (Moradi <i>et al.</i>, 2018), which are similar peak with this present study.</p>
6.	There were no distinct results between the two selected methods of study with the combined methods of study - justification is to be required.	<p>Distinct results between two selected methods of study with the combined methods of study has been added in the manuscript. Please see the red highlighted of:</p> <p>Results and discussion Page 9, Line 23- Line 29: In this study, combining the derived fraction obtained from HPSEC-OCD and -UVD with peak-fitting techniques and the derived component from F-EEM with FRI analysis, on the same sample simultaneously, gave a reinforced information and had confirmed the presence of humic substances-like (Peak B)/humic acid-like (Region 5) as the main organic in nitrification processes. The study indicates that a combination of HPSEC-OCD/UVD and FEEM can be used to determine the characteristics of organic matter and water quality change during nitrification.</p>

7.	There was no informative results (values) presented in the abstract – include them.	Informative results (values) in term of percentage area has been added in the abstract. Please see the red highlighted of: Abstract Page 1 Line 9 – Line 12, and Line 17. According to chromatography with carbon and ultraviolet detector , the main characteristics of organic matter shifted from building blocks aromatic compounds with percentage peak area of carbon/ultraviolet detector: 31%/53% to 14%/27.5% to humic-like substances with percentage peak area of carbon/ultraviolet detector 21%/17% to 27%/46.5% during nitrification. Those former compounds are biodegradable as well as properties of microbial products released during substrate utilization and endogenous phase, which are mainly identified as humic-like substances, thus underwent further biodegradation. However, there was significant change in the fluorophores organic fractions, which exhibited humic acid-like with percentage fluorescence regional index area 53% into 68% , as shown by fluorescence excitation-emission matrix analysis.
----	---	---

Reviewer # 2:

Query	Details reviewing required	Author's response
1.	Abstract; do not use abbreviation "FEEM"	Abbreviation FEEM has been deleted in the abstract. Please see the revised abstract.
2.	Abstract; add more data for the results, Abstracts should be able to stand alone.	<p>More data for the results has been added in abstract. Please see the red highlighted of:</p> <p>Abstract Page 1 Line 9 – Line 12, and Line 17.</p> <p>According to chromatography with carbon and ultraviolet detector, the main characteristics of organic matter shifted from building blocks aromatic compounds with percentage peak area of carbon/ultraviolet detector: 31%/53% to 14%/27.5% to humic-like substances with percentage peak area of carbon/ultraviolet detector 21%/17% to 27%/46.5% during nitrification. Those former compounds are biodegradable as well as properties of microbial products released during substrate utilization and endogenous phase, which are mainly identified as humic-like substances, thus underwent further biodegradation. However, there was significant change in the fluorophores organic fractions, which exhibited humic acid-like with percentage fluorescence regional index area 53% into 68%, as shown by fluorescence excitation-emission matrix analysis.</p>
3.	Complete the sentence at the end of introduction	<p>The sentence at the end of introduction has been completed. Please see the red highlighted of:</p> <p>Introduction Page 2, Line 48- Line 49.</p> <p>This study has been carried out in laboratory batch scale at Department of Environmental Engineering, National Cheng Kung University, Taiwan in 2016.</p>
4.	Add original references to each section of materials and methods	<p>Original references have been added to section of material and methods, references as well.</p> <p>Please see the red highlighted of :</p> <p>Material and methods Page 3, Line 21, Line 31, Line 36.</p> <p>(Liu 2013), (Murphy <i>et al.</i>, 2013), (Chow <i>et al.</i>, 2008).</p>