

DAFTAR PUSTAKA

- Afiadi, F. (2015) 'Model Estimasi Variasi Spasial Seismisitas Sebagai Prekursor Gempa Bumi Kuat Menggunakan Extreme Learning Machine (ELM)', in *TESIS Program Studi Instrumentasi dan Kontrol*. ITB.
- Demertzis, K. and Iliadis, L. (2016) 'Bio-inspired hybrid intelligent method for detecting android malware', in *Knowledge, Information and Creativity Support Systems*. Springer, pp. 289–304.
- Gutenberg B. dan Richter, C. F. (1944) 'Frequency of earthquakes in California, Bull.', *Seismol. Soc. Am.*, 34.
- Hamilton, W. B. (1979) 'Tectonics of the Indonesian region', in *US Geol. Surv. Prof. Pap.*, 1078, 345. doi: 10.3133/pp1078.
- Huang, G.-B. *et al.* (2008) 'Incremental extreme learning machine with fully complex hidden nodes', *Neurocomputing*. Elsevier, 71(4), pp. 576–583.
- Huang, G.-B., Zhu, Q.-Y. and Siew, C.-K. (2004) 'Extreme learning machine: a new learning scheme of feedforward neural networks', in *Neural Networks, 2004. Proceedings. 2004 IEEE International Joint Conference on*, pp. 985–990.
- Huang, G.-B., Zhu, Q.-Y. and Siew, C.-K. (2006) 'Extreme learning machine: theory and applications', *Neurocomputing*. Elsevier, 70(1), pp. 489–501.
- Joelianto, E., Widiyantoro, S. and Ichsan, M. (2008) 'Time series estimation on earthquake events using ANFIS with mapping function', *International Journal of Artificial IntelligenceTM*, 3(A09), pp. 37–63.
- Kumar, S. (2012) 'Seismicity In The Nw Himalaya, India: Fractal Dimension, B-

- Value Mapping And Temporal Variations For Hazard Evaluation’, *Geoscience Research*, 3, pp. 83–87.
- Kundu, B. dan G. V. K. (2011) ‘Slab Detachment Of Subducted Indo-Australian Plate Beneath Sunda Arc, Indonesia’, *J. Earth Syst. Sci.*, 120(193), pp. 193–204. doi: <https://doi.org/10.1007/s12040-011-0056-7>.
- Pratiwi Rohadi, S. N. A. N. (2012) ‘Analisis korelasi variasi spasial dan temporal b-value terhadap stress, seismisitas, dan tektonik di pulau bali, lombok, dan Sumbawa’, *Jurnal. Geofisika*, 13(2/2012).
- Rahmat, B., Afiadi, F. and Joelianto, E. (2018) ‘Earthquake Prediction System using Neuro-Fuzzy and Extreme Learning Machine BT - International Conference on Science and Technology (ICST 2018)’, in. Atlantis Press. doi: <https://doi.org/10.2991/icst-18.2018.95>.
- Rohadi, S. (2009) ‘Studi Seismotektonik Sebagai Indikator Potensi Gempa bumi Di Wilayah Indonesia’, *Jurnal Meteorologi Dan Geofisika*, 10, pp. 111–120.
- Schorlemmer D., S. W. and Wyss, M. (2004) ‘Earthquake statistics at Parkfield, Stationarity of b-values’, *J. of Geophys. Res.*, 109(B12307). doi: [doi: 10.1029/2004-JB003234](https://doi.org/10.1029/2004-JB003234).
- Shi Y. and Bolt, B. A. (1982) ‘The standard error of the magnitude-frequency b value’, *Bull. Seismol. Soc. Am.*, 72, pp. 1677–1687.
- Sieh K., N. D. (2000) ‘Neotectonics of the Sumatran Fault Indonesia’, *Journal of Geophysical Research*, 105, pp. 28295–28326.
- Sompotan, A. *et al.* (2015) ‘Analysis of Ionospheric Precursor of Earthquake using GIM-TEC, Kriging and Neural Network’, *Asian Journal of Earth Sciences*, 8, pp. 32–44. doi: [10.3923/ajes.2015.32.44](https://doi.org/10.3923/ajes.2015.32.44).

- Sunardi, B. (2009) 'Analisa Fraktal Dan Rasio Slip Daerah Bali-Ntb Berdasarkan Pemetaan Variasi Parameter Tektonik', *Jurnal Meteorologi dan Geofisika*, 10, pp. 58–65.
- Ustu, T. (1965) 'A method in determining the value of b in a formula $\log n = a - bM$ showing the magnitude frequency for earthquakes', *Geophys. Bull. Hokkaido Univ.*, 13, pp. 99–103.
- Wiemer S. and Wyss, M. (2002) 'Mapping spatial variability of the frequency-magnitude distribution of earthquakes', *Adv. Geophys.*, 45, pp. 259–302.